

D-Branes, AdS/CFT Correspondence, and T-Duality

Yuri Malyuta

Institute for Nuclear Research

National Academy of Sciences of Ukraine

252022 Kiev, Ukraine

e-mail: interdep@kinr.kiev.ua

Abstract

The D -brane spectra in the Type IIB string theory compactified on $AdS_{p+2} \times S^{8-p}$ are computed using the K-theory approach. The results signal the existence of the mirror-symmetry-analogue for D -branes, analogous to that risen in the context of derived categories.

Keywords: D-branes, String Duality, Superstring Vacua.

1 Introduction

D -branes play a significant role in superstrings and superconformal field theories. Two of the most outstanding developments in this direction have been achieved:

1. The generalized AdS/CFT correspondence [1], which relates the superconformal field theory on Dp -branes placed at the orbifold singularity and the Type IIB string theory compactified on $AdS_{p+2} \times H^{8-p}$;
2. The K-theory approach to D -brane charges [2, 3, 4], which identifies D -brane charges with elements of Grothendieck K-groups [5, 6, 7] of horizon manifolds.

In the present paper we use K-theory to compute the D -brane spectra in the Type IIB string theory compactified on $AdS_{p+2} \times S^{8-p}$.

2 D -brane spectra

Let us consider the fibre bundle

$$\begin{array}{ccc} S^{8-p} & \rightarrow & B^{9-p} \\ & \downarrow & \\ & & B^{9-p}/S^{8-p} \end{array}$$

The K-groups characterizing this bundle are related by the exact hexagon [7]

$$\begin{array}{ccc}
& \widetilde{K}(B^{9-p}/S^{8-p}) & \rightarrow & \widetilde{K}(B^{9-p}) \\
& \nearrow \delta & & \searrow \\
\widetilde{K}(SS^{8-p}) & & & \widetilde{K}(S^{8-p}) \\
& \nwarrow & & \swarrow \delta \\
& \widetilde{K}(SB^{9-p}) & \leftarrow & \widetilde{K}(S(B^{9-p}/S^{8-p}))
\end{array}$$

where δ is the coboundary homomorphism. This hexagon is the counterpart of the generalized AdS/CFT correspondence (cf. [3]).

Since

$$\widetilde{K}(B^{9-p}) = \widetilde{K}(SB^{9-p}) = 0 ,$$

the hexagon splits into the exact sequences

$$0 \rightarrow \widetilde{K}(SS^{8-p}) \xrightarrow{\delta} \widetilde{K}(B^{9-p}/S^{8-p}) \rightarrow 0 \quad (1)$$

$$0 \rightarrow \widetilde{K}(S^{8-p}) \xrightarrow{\delta} \widetilde{K}(S(B^{9-p}/S^{8-p})) \rightarrow 0 \quad (2)$$

The sequences (1) and (2) are related by T-duality.

The group $\widetilde{K}(SS^{8-p})$ from (1) reproduces the D -brane spectrum

Table 1

Dp	$D9$	$D8$	$D7$	$D6$	$D5$	$D4$	$D3$	$D2$	$D1$	$D0$	$D(-1)$
S^{9-p}	S^0	S^1	S^2	S^3	S^4	S^5	S^6	S^7	S^8	S^9	S^{10}
$\widetilde{K}(S^{9-p})$	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$

which coincides with the known result in Type IIB theory [8].

The group $\widetilde{K}(S^{8-p})$ from (2) reproduces the D -brane spectrum

Table 2

Dp	$D9$	$D8$	$D7$	$D6$	$D5$	$D4$	$D3$	$D2$	$D1$	$D0$	$D(-1)$
S^{8-p}	S^{-1}	S^0	S^1	S^2	S^3	S^4	S^5	S^6	S^7	S^8	S^9
$\widetilde{K}(S^{8-p})$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0	$\mathbb{Z}$	0

which signals the existence of the mirror-symmetry-analogue for branes, analogous to that rised in the context of derived categories [4].

3 Vacuum manifold

Using standard definitions [7], we obtain

$$\widetilde{K}(SS^{8-p}) = \pi_{9-p}(BU) ,$$

$$\widetilde{K}(S^{8-p}) = \pi_{8-p}(BU) ,$$

where BU is the inductive limit of the manifold

$$U(2N)/U(N) \times U(N) \tag{3}$$

The vacuum manifold (3) has the following interpretation in terms of D -branes [9]. When $2N$ coinciding branes are separated to form two parallel stacks of N coinciding branes, their gauge symmetry $U(2N)$ is spontaneously broken to $U(N) \times U(N)$. This situation generically allows for the existence of topological solitons.

4 Acknowledgements

I would like to thank S. Gukov for stimulating discussions. It is a pleasure to thank E. Witten for his attention to this work.

References

- [1] D.R. Morrison and M.R. Plesser, *Non-Spherical Horizons, I*, hep-th/9810201.
- [2] E. Witten, *D-Branes and K-Theory*, hep-th/9810188.
- [3] S. Gukov, *K-Theory, Reality, and Orientifolds*, hep-th/9901042.
- [4] E. Sharpe, *D-Branes, Derived Categories, and Grothendieck Groups*, hep-th/9902116.
- [5] M.F. Atiyah, *K-Theory* (Benjamin, New York 1967).
- [6] M. Karoubi, *K-Theory. An Introduction* (Springer, Berlin 1978).
- [7] D. Husemoller, *Fibre Bundles* (McGraw-Hill, New York 1966).
- [8] K. Olsen and R.J. Szabo, *Constructing D-Branes from K-Theory*, hep-th/9907140.
- [9] G.W. Semenoff and K. Zarembo, *Solitons on Branes*, hep-th/9903140.